Limonia flavipes L. macrostigma L. maculipennis L. nigropunctata A strikingly marked striped brown species. Spring A brown species with unmarked wings, and quite A very striking black and orange species with An orange species with a dark thoracic stripe, striped orange and black legs. Flies in April and to summer in woodland, particularly calcareous different to other woodland Limonia. Found by though sometimes a darker with an all black May. Associated with hedgerows and a limited woodland. Has a fingerprint mark on the side and woodland streams, and wet woodlands. May to thorax. Found in woodland. Striking glossy species range of other sites. Ecology uncertain. rear of the thorax (microscope needed to view it October. Vein Sc at or just past base of vein Rs. - wing with 3 spots and dark wing edge. Legs can Restricted mostly to the south-east of England. be dark-tipped or nearly all dark. April – June. clearly).

L. nubeculosa	L. phragmitidis	L. stigma	L. trivittata
A very common species of shady places potentially found throughout the year. Wing has unique patterning and legs have two pales stripes on the femur.	A common spring woodland species. Grey head, single stripe at the front of the thorax, and three wing spots.	An uncommon summer woodland species. Yellow head, single stripe at the front of the thorax, and three wing spots.	A local late spring species of damp woodland, often dingle woodlands. Three pale barely discernible wing spots, grey head, and three thoracic stripes.

Crib to UK woodland *Limonia* **sp. craneflies** – one non-woodland species omitted (*Limonia dilutior* found on heathland and moorland). For educational purposes only – not for publication. UK Cranefly Recording Scheme – Pete Boardman, 2020.