





















































| Taxon / Status | Habitat | Head | Thorax | Abdomen | Wing | cerci |
|---|--|---|---|--|---|-------|
| <i>N. analis</i> (Local) Widespread | Usually near rivers & streams – sandy substrates |  |  |  |  | blunt |
| <i>N. appendiculata</i> (Very common) | Gardens, all but the wettest grassland, verges, etc. |  |  |  |  | sharp |
| <i>N. cornicina</i> (Common) Commoner in the southern drier areas | Grassland, including dry grassland. |  |  |  |  | sharp |
| <i>N. crocata</i> (Restricted) More frequent in south & midlands | Sandy soils, quarries, peaty soils, heaths. |  |  |  |  | blunt |
| <i>N. dorsalis</i> (Local) Commoner in the north & west | Sandy river banks. Mostly northern |  |  |  |  | blunt |
| <i>N. flavescens</i> (Very common) | Rough grassland, fields, coastal grassland. |  |  |  |  | sharp |
| <i>N. flavipalpis</i> (Common) Except far north | Woodland, woodland edge, hedgerows |  |  |  |  | sharp |

| | | | | | | |
|--|---------------------------------|---|---|--|---|---------------------|
| <i>N. guestfalica</i> (Local) Widespread | On sandy soils. |  |  |  |  | sharp |
| <i>N. lunulicornis</i> (Restricted) Mostly northern & western | Rivers in the uplands & western |  |  |  |  | blunt black edge |
| <i>N. quadrifaria</i> (Very common) | Woodland edge, gardens. |  |  |  |  | blunt |
| <i>N. quadristriata</i> (Restricted) Mostly coastal, west and south | Coastal |  |  |  |  | blunt |
| <i>N. scurra</i> (Local) Widespread | Sandy habitats |  |  |  |  | blunt |
| <i>N. submaculosa</i> (Restricted) Mostly coastal & northern river banks, Brecks. | Coastal dunes and inland sands |  |  |  |  | sharp |

UK 'tiger craneflies' *Nephrotoma* (Diptera: Tipulidae) species crib (minus two very restricted rarities – *N. aculeata* (Scottish Highlands only) & *N. sullingtonensis* (one Sussex heathland)). Note – if unsure please key out using Stubbs & Kramer, 2016 or Boardman, 2016 for more info.

FOR EDUCATIONAL USE ONLY – NOT FOR PUBLICATION.

Pete Boardman UK Cranefly Recording Scheme 2020.